



Learn and Live..

[Print this Page for Your Records](#)[Close Window](#)**Control/Tracking Number:** 07-SS-A-12233-AHA**Activity:** Abstract**Current Date/Time:** 5/29/2007 5:27:12 PM**Persistent Gender Bias in the Use of Percutaneous Coronary Intervention at the Acute Stage of MI from 1995 to 2005: Results from the French USIK, USIC 2000 and FAST-MI Registries**

Author Block: Nicolas DANCHIN, HEGP, Paris, France; Tabassome SIMON, CHU St Antoine, Paris, France; Helene ELTCHANINOFF, CHU Rouen, Rouen, France; Martine GILARD, CHU Brest, Brest, France; Ioana OLTEAN, CH Cambrai, Cambrai, France; Florence LECLERCQ, CHU Montpellier, Montpellier, France; Geneviève MULAK, Jean-Pierre CAMBOU, SFC, Paris, France; for the USIK 1995, USIC 2000 and FAST-MI investigators

Abstract: Percutaneous coronary intervention (PCI) is an essential step in the management of patients with STEMI or NSTEMI. Previous data have evidenced a gender bias in the use of PCI in people admitted for AMI.

Aim: to assess whether PCI was used differently in men and women with AMI and if there have been changes in this regard over the last 10 years.

Methods: USIK 1995, USIC 2000 and FAST-MI are 3 nationwide French registries conducted 5 years apart, using a similar methodology in patients admitted to CCU over a one-month period. For the present analysis, all 7528 patients admitted for AMI (STEMI and NSTEMI) < 48 hours from symptom onset were included.

Results: 2152 patients were included in 1995, 2317 in 2000 and 3059 in 2005. Women represented 29%, 27% and 32% of the populations, respectively in the 3 periods. Overall, women were 10 years older than men (no difference between the 3 periods); PCI was used in 53% of men and 39% of women ($p < 0.001$). Use of PCI during the initial hospital stay increased from 1995 to 2005, but the difference between men and women persisted (1995: 12% v. 20%; 2000: 45% v. 62%; 2005: 52% v. 69%) ($p < 0.001$ at all time points). In the whole population, female gender was an independent predictor of a lower use of reperfusion therapy (OR: 0.70; 95% CI: 0.62-0.79, $p < 0.001$), together with older age, history of peripheral arterial disease, history of congestive heart failure, and earlier time period (1995 v 2000 v 2005) (logistic regression analysis). In each of the 3 periods, female gender remained associated with lower use of reperfusion, with odds ratios of similar magnitude (1995: OR: 0.54, 95% CI: 0.41-0.71; 2000: OR 0.50, 95% CI: 0.41-0.60; 2005: OR: 0.48, 95% CI: 0.41-0.56). A striking decrease in 30-day mortality was observed in women (1995: 21%, 2000: 12%, 2005: 8.5%). This decrease, however, was found predominantly in women who underwent PCI (14%, 6% and 4%, respectively), compared with women without PCI (22%, 17%, 14%, respectively). **Conclusion:** In spite of a much higher use of PCI over the past 10 years, a definite gender bias persists. This finding is all the more preoccupying as the reduction in early mortality over this period of time has been observed mainly in women with PCI.

Author Disclosure Block: N. Danchin, The FAST-MI registry was supported by a grant from Pfizer and Servier and USIK1995 and USIC 2000 were supported by Aventis, Significant,2. Research Grant; T. Simon, None; H. Eltchaninoff, None; M. Gilard, None; I. Oltean, None; F. Leclercq, None; G. Mulak, None; J. Cambou, None.

AHA Awards (Complete):**No, I am not interested in AHA Early Career Investigator Awards or Travel Stipends : True****Category (Complete):** Unstable Angina, NSTEMI and STEMI: Prognosis and Pharmacologic Therapy (Non-Catheter Based) (CLCD)

Keyword (Complete): Acute Coronary Syndromes ; Outcomes ; Angioplasty ; Gender

Presentation Preference (Complete): All Formats

Additional Info (Complete):

: A. Scientific Sessions Web site (scientificsessions.org)

Payment (Complete): Your credit card order has been processed on Tuesday 29 May 2007 at 5:25 PM.

Status: Complete

American Heart Association

7272 Greenville Avenue

Dallas, Texas 75231

Technical Support: support@abstractsonline.com or 1-217-398-1792

Powered by OASIS, The Online Abstract Submission and Invitation System SM

© 1996 - 2007 Coe-Truman Technologies, Inc. All rights reserved.